

PUBBLICAZIONI SU RIVISTE INTERNAZIONALI

Continuous i.v. infusion of remifentanil and intraosseous lidocaine provide better analgesia than intraosseous lidocaine alone in percutaneous vertebroplasty of osteoporotic fractures.

Dauri M, Coniglione F, Faria S, Fiori R, Frunzo F, Massari F, Simonetti G, Sabato AF, Masala S.

Br J Anaesth. 2009 Dec;103(6):901-2. No abstract available.
PMID: 19918028 [PubMed - indexed for MEDLINE]

White matter reduced streamline coherence in young men with autism and mental retardation.

Pardini M, Garaci FG, Bonzano L, Roccatagliata L, Palmieri MG, Pompili E, Coniglione F, Krueger F, Ludovici A, Floris R, Benassi F, Emberti Gialloreti L. Eur J Neurol. 2009 Nov;16(11):1185-90. Epub 2009 Jun 15.
PMID: 19538216 [PubMed - indexed for MEDLINE]

Continuous femoral nerve block provides superior analgesia compared with continuous intra-articular and wound infusion after anterior cruciate ligament reconstruction.

Dauri M, Fabbi E, Mariani P, Faria S, Carpenedo R, Sidiropoulou T, Coniglione F, Silvi MB, Sabato AF.

Reg Anesth Pain Med. 2009 Mar-Apr;34(2):95-9.
PMID: 19282706 [PubMed - indexed for MEDLINE]

A new technique for continuous intercostal-intrapleural analgesia in videothoracoscopic surgery.

Forcella D, Pompeo E, Coniglione F, Gatti A, Mineo TC.

J Thorac Cardiovasc Surg. 2009 Jan;137(1):e48-9. Epub 2008 Aug 27. No abstract available.
PMID: 19154885 [PubMed - indexed for MEDLINE]

Comparison of epidural, continuous femoral block and intraarticular analgesia after anterior cruciate ligament reconstruction.

Dauri M, Polzoni M, Fabbi E, Sidiropoulou T, Servetti S, Coniglione F, Mariani P, Sabato AF.

Acta Anaesthesiol Scand. 2003 Jan;47(1):20-5.
PMID: 12492792 [PubMed - indexed for MEDLINE]

